

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

THIS PAGE BLANK (USPTO)

PCT/IL 99 / 00541

25 NOVEMBER 1999

PA 170382

REC'D 06 DEC 1999

TO PCT

THE UNITED STATES OF AMERICA

TO ALL TO WHOM THESE PRESENTS SHALL COME:

UNITED STATES DEPARTMENT OF COMMERCE

United States Patent and Trademark Office

November 04, 1999

THIS IS TO CERTIFY THAT ANNEXED HERETO IS A TRUE COPY FROM THE RECORDS OF THE UNITED STATES PATENT AND TRADEMARK OFFICE OF THOSE PAPERS OF THE BELOW IDENTIFIED PATENT APPLICATION THAT MET THE REQUIREMENTS TO BE GRANTED A FILING DATE UNDER 35 USC 111.

APPLICATION NUMBER: 60/104,118

FILING DATE: October 13, 1998

09/856429

PRIORITY DOCUMENT
SUBMITTED OR TRANSMITTED IN
COMPLIANCE WITH
RULE 17.1(a) OR (b)



By Authority of the
COMMISSIONER OF PATENTS AND TRADEMARKS

P. SWAIN

Certifying Officer

November 04, 1999

CHRISTINA
PSF BOX: 207

This is to inform you that the document requested in your order dated 11-02-1999 is a true reproduction of the official office record copy of that document:

60104118 CERTIFIED PAT APP AS FILED-EXPEDITE-PSF

- ☐ The enclosed Patent Application as Filed is a reproduction of the application as originally filed and has been recorded using high quality scanning or microfilm equipment. Copies of page/papers that were not scannable have not been included, nor have pages/papers received after the original filing date. Copies of these pages/papers may be ordered separately.
- ☒ The enclosed document is a reproduction of the best available source of the official office record copy of that document.

If you have any questions or need additional information, please contact our Customer Service Department.

Mailing Address:

U.S. Patent and Trademark Office
Office of Public Records, Customer Service
Crystal Gateway 4, Suite 300
Washington DC 20231

Delivery Address:

U.S. Patent and Trademark Office
Office of Public Records, Customer Service
1213 Jefferson Davis Highway, Suite 300
Arlington VA 22202

For faster processing of new orders, please specify as appropriate:

Box 9 (Copy Sales) for Uncertified copies, or Box 10 for Certified copies of PTO Documents

Voice: (703) 308-9726 Fax: (703) 308-7048 E-Mail: PTCS@USPTO.GOV or
Certdiv@USPTO.GOV

Ref:PS 170382

PROVISIONAL APPLICATION COVER SHEET

This is a request for filing a PROVISIONAL APPLICATION under 37 CFR 1.53(b)(2).

Docket Number: 1772/44761PV

Type a plus sign (+) inside this box -

10/13/98

JC408 U.S. PTO

INVENTOR(S)/APPLICANT(S)			
LAST NAME	FIRST NAME	MIDDLE INITIAL	RESIDENCE (CITY AND EITHER STATE OR FOREIGN COUNTRY)
Nussinovitch Kampf	A. N.	-- --	Jerusalem, Israel Jerusalem, Israel
TITLE OF THE INVENTION (280 characters max)			
Hydrocolloid Coating of Embryos			
CORRESPONDENCE ADDRESS			
Richard R. Diefendorf Evanson, McKeown, Edwards & Lenahan, P.L.L.C. 1200 G. Street, N.W., Suite 700 Washington, D.C. 2005 USA			
ENCLOSED APPLICATION PARTS (check all that apply)			
<input checked="" type="checkbox"/> SPECIFICATION	NUMBER OF PAGES 4	<input type="checkbox"/> SMALL ENTITY STATEMENT	
<input checked="" type="checkbox"/> DRAWINGS	NUMBER OF SHEETS 2	<input type="checkbox"/> OTHER (specify) _____	
METHOD OF PAYMENT (check one)			
<input checked="" type="checkbox"/> A check or money order is enclosed to cover the Provisional filing fees	PROVISIONAL FILING FEE AMOUNT (\$) \$150		
<input type="checkbox"/> The Commissioner is hereby authorized to charge filing fees and credit Deposit Account Number: 05-1323			

JC541 U.S. PTO

60/101116

10/13/98

The invention was made by an agency of the United States Government or under a contract with an agency of the United States Government.

☒ No.

☐ Yes, the name of the U.S. Government agency and the Government contract number are:

Respectfully submitted,

SIGNATURE Richard R. Diefendorf
TYPED or PRINTED NAME Richard R. Diefendorf

Date October 13, 1998
REGISTRATION NO. 32,390
(if appropriate)

☐ Additional inventors are being named on separately numbered sheets attached hereto

PROVISIONAL APPLICATION FILING ONLY

Hydrocolloid coating of embryos

A. Nussinovitch and N. Kampf

Institute of Biochemistry, Food Science and Nutrition
Faculty of Agricultural, Food and Environmental Quality Sciences
The Hebrew University of Jerusalem

Idea

Coating embryos by hydrocolloidal thin films to achieve:

- a) Postpone hatching and extended survival rates.
- b) Protection from microbial contamination
- c) Protection from hazardous materials produced or introduced into the media.
- d) As an inhibitor against damages occurred during freezing and thawing.

Example 1

X. laevis eggs were coated immediately after squeezing and fertilization by a thin layer ($\sim 50 \mu$) of film based on three different types of alginates varying in their mannuronic/guluronic ratios (Fig. 1a and b). The alginate was cross-linked either by Ca or Ba ions at three different concentrations. The development, survival and hatching of these embryos and the swelling of their natural jelly coats or hydrocolloid coating were studied during 7 days, while embryos were maintained either in flowing aerated water at a ratio of 85 ml per embryo or at a very diminished ratio of 0.6 ml sterile or non-sterile modified Marc Ringer's solution per embryo. All experiments were conducted in triplicates at $20 \pm 1^\circ \text{C}$. Oxygen was monitored continuously.

The coatings succeeded in postponing hatching in ca. 60 h in flowing aerated water at a ratio of 85 ml per embryo. However, the survival prospects diminished. Calcium as cross-linking agent was found to a better contribution.

However, major advantages of the coating were observed when the ratio between the embryos and the surrounding medium was maintained at its minimal value in non-sterile conditions, perhaps due to film resistance to diffusion. Here 0.25% barium and 0.25% calcium as cross-linking agents of the alginates gave the best results. In the studied systems, the coating seemed to postpone the hatching of the embryos. The difference in hatching time between the blank and the coated embryos was 30 to 60 h. In addition, the coating served as a barrier to microbial contamination and thus improved survival prospects. The number of microorganisms, counted directly at the medium after 5 days was 10^3 to 10^6 CPU, depends on conditions, medium temperatures and ratio between volume of medium and embryos number.

Example 2

Three different kinds of alginate with different glucuronic (G) to mannuronic (M) acid ratios have been tried in coating the embryos. It is observed that the lower M to G ratio achieved better results with % of hatching. When this ratio was higher less penetration of high molecular weight compounds occur. Thus choosing the appropriate combination will determine the success of the coating.

Example 3

X. laevis eggs were coated immediately after squeezing and fertilization by a thin layer of films based on LMP (Low Methoxy Pectin), λ -carrageenan and

α-carrageenan. The LMP was cross-linked either by Ca or Ba ions at different concentrations (other cross-linking ions are also possible). The λ-carrageenan and κ-carrageenan were cross-linked by Ca and potassium ions respectively at different concentrations. The development, survival and hatching of these embryos were studied during 7 days, while embryos were maintained at a ratio of 0.6 ml non-sterile modified Marc Ringer's solution per embryo. All experiments were conducted in triplicates at $20 \pm 1^\circ\text{C}$. For the λ-carrageenan and κ-carrageenan coated embryos, a higher survival rate than the non-coated embryos (calculated as a percent of the total hatching embryos) was observed. 1% LMP and 1% alginate were less effective. In fact, all of these coatings improved the survival rates under experiment conditions. The major advantages of the coating were observed when the ratio between the embryos and the surrounding medium was maintained at its minimal value in non-sterile conditions, perhaps due to film resistance to diffusion. In these systems, the coating seemed to postpone the hatching of the embryos. In addition, the coating served as a barrier to microbial contamination and thus improved survival prospects.

Example 4

Same as proposed for examples 1 and 3. The coated embryos were frozen by Planer cryo 10 at a rate $< 1^\circ\text{C}/\text{min}$ up to a temp of -7°C and then at a freezing rate $> 10^\circ\text{C}/\text{min}$ to -50°C . The coated frozen embryos were transferred to liquid nitrogen. The embryos were kept at each step for a few minutes for temperature stabilization, completing of crystallization and to permit majority of water to leave the cell. The percentages of embryo survival after one cycle of freezing and thawing were higher (in ~5 to 30%) than what that observed for the non-coated embryos. It is proposed that this result is the outcome of two


mechanisms. In the first, the coating served as a mechanical membrane and eliminates in part the penetration of the embryo by medium icicles. The second mechanism is probably the outcome of disturbance to ice to crystalize during the freezing.

Figs. Legends

Fig. 1a and b: embryo after fertilization and in advanced stage coated by alginate film.

Fig. 2a and b: emergence of embryos from the hydrocolloid coating

United States Patent & Trademark Office
Office of Initial Patent Examination -- Scanning Division



Application deficiencies were found during scanning:

☐ Page(s) _____ of _____ were not present
for scanning. (Document title)

☐ Page(s) _____ of _____ were not present
for scanning. (Document title)

☐ Scanned copy is best available.

NO DECLARATION

Fig. 1a



Fig. 1b



Fig. 2a




Fig. 2b

